

9100154

THIE UNITHED STATES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

pioneer Gi-Bred International, Inc.

Colleceas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, importing it, or exporting it, or using it in producing a hybrid or different ty therefrom, to the extent provided by the Plant Variety Protection Act 1542, as amended, 7 u.s.c. 2321 et seq.)

ALFALFA

152521

In Lestimony Whereot, I have hereunto set my hand and caused the seal of the Plant Tariety Protection Office to be affixed at the City of Washington, D.C.

this 30th day of April in the year of our Lord one thousand nine Jundred and ninety-three.

Clika Ess

dure

Kenneth Hours

Commissioner

Plant Variety Protection Office Agricultural Marketing Service Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250, and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Expires 1/31/91

| U.S. DEPARTMENT OF AGRICU AGRICULTURAL MARKETING SI | ULTURE SERVICE | | Application is required i | in order to |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|
| APPLICATION FOR PLANT VARIETY P | ROTECTION | CERTIFICATE | determine if a plant variet certificate is to be issued (7 Information is held conti- certificate is issued (7 U.S.C. | y profection USC, 2421) dential until |
| 1. NAME OF APPLICANT(S) (as it is to appear on the Certificate) | | 2. TEMPORARY DESIGNATION (| DR 3. VARIETY NAME | ···· |
| Pioneer Hi-Bred International, Inc. | | EXPERIMENTAL NO. XAM93 | 5252 | _ |
| 4 ADDRESS (street and no. or R.F.D. no., city, state, and ZIP) | | 5. PHONE (Include area code) | FOR OFFICIAL USE | ONLY: |
| 7305 N. W. 62nd Avenue, P.O. Box 287 | | • | PVPO NUMBER | |
| Johnston, IA 50131 | | 515-270-3340 | 2,22 | |
| | | | 91001 | 54 |
| | | · · | F Date | - 1001 |
| | AMILY NAME (Bolanic | al) | March 2 | 16222 |
| Medicago sativa | eguminosae | | N G A M | и. 🔲 Р М |
| 8 CROP KIND NAME (Common Name) | 9. 0 | ATE OF DETERMINATION | F Filing and Examinate | on Fee. |
| Alfalfa | | August, 1989 | [] 2/30· | |
| 10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGANIZATION | ION (Corporation, parti | ership, association, etc.) | | 1991 |
| Corporation | | | C Certificate Fee: | 177 F |
| 11. IF INCORPORATED, GIVE STATE OF INCORPORATION | 12. DA | TE OF INCORPORATION | - \$ \$250.00 | |
| Iowa | | 1926 | V Date | 20.0 |
| 13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE | E IN THIS APPLICATIO | N AND RECEIVE ALL PAPERS | 6 apr. 15,19 | 73 |
| Mary Helen Mitchell, 700 Capital Square, John Hintze, 700 Capital Square, 400 Loc John Hintze, 700 Capital Square, 400 Loc IIII CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow INS a Exhibit A, Origin and Breeding History of the Variety. b. Exhibit B, Novelty Statement. c. Exhibit C, Objective Description of Variety. d Exhibit D, Additional Description of Variety. e Exhibit E, Statement of the Basis of Applicant's Ownership. I Seed Sample (2,500 viable untreated seeds) Date Seed Sample (2,500 | cust Street STRUCTIONS on revers ole mailed to Plant V arer of the United Sta VARIETY NAME ONLY 17. IF "YES" TO 17. IF "YES" TO 18. THE U.S.? | ariety Protection Office Mailes." AS A CLASS OF CERTIFIED SEED." skip to item 18 below) ITEM 16, WHICH CLASSES OF PE | 50309 s code): rch 13, 1991 Properties the Plant Var. RODUCTION BEYOND BREEDER SEE | |
| ™ NO | · violation one data | · · · · · · · · · · · · · · · · · · · | | |
| 20 The applicant(s) declare(s) that a viable sample of basic seeds of request in accordance with such regulations as may be applicable. The undersigned applicant(s) is (are) the owner(s) of this sexus uniform, and stable as required in section 41, and is entitled to p. Applicant(s) is (are) informed that false representation herein ca | d for U.S.A Ithis variety will e. ally reproduced reprotection under the | Spring, 1992 be furnished with the appli ovel plant variety, and be e provisions of section 42 of ction and result in penalties | lieve(s) that the variety is di the Plant Variety Protection 2 | iclinet |
| PIONEER HI-BRED INTERNATIONAL, INC. | | | | |
| BY Alleam TM Alongward | l l | TLE C, Department of Breeding | 3-12-9. | / |

EXHIBIT A

ORIGIN AND BREEDING HISTORY OF THE VARIETY

152521

5252 is a synthetic variety with 197 parent plants originating from an experimental line tracing to the varieties 5373 and Surpass. Parent plants were selected through phenotypic recurrent selection for resistance to Phytophthora root rot and/or Aphanomyces root rot. Germplasm sources are M. falcata (8%), Ladak (8%), M. varia (23%), Turkistan (10%), Flemish (42%), Chilean (7%), and Peruvian (2%).

During seed multiplication no variates beyond the limits defined under Exhibit C have been found. Multiplication procedures will insure that seed being sold as 5252 will not be shifted in characteristics beyond presently acceptable limits for alfalfa varieties. Syn 1 seed harvested from individual plants in "cage isolation" in 1989 is considered breeder seed.

It is confirmed that 5252 meets presently acceptable levels for uniformity for alfalfa varieties.

EXHIBIT B

NOVELTY STATEMENT

152521

5252 most closely resembles the variety 'Arrow'. 5252 differs from Arrow in anthracnose resistance, being classified as having resistance, while Arrow has moderate resistance to the disease.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT GRADE & BEFO DIVISION PLANT SARGITY PROTECTION OFFICE BEETSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

| | | ALFALFA | (Medicago sativa sei | nsu Gunn et al.) | | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|---------------------------------------|---------------------------------------------------------------------------|----------------------------------|----------------------------------------|---------------------------------------|
| NAME OF APPLICANT(S) | | | TEMPORARY | DESIGNATION | VARIETY NAME | | |
| Pioneer Hi-Bred Into | ernational, | Inc. | хам9 | 3 | 5252 | | |
| ADDRESS (Street and No., or R.F.D. | No., City, State, and Zi | p Code) | | | FOR OFFICIAL USE ONLY | | |
| 7305 N. W. 62nd Aver | nue, P. O. : | Box 287 | | | PVPO NUMBER | | |
| Johnston, IA 501 | 31 | | | | | 9100154 | <u> </u> |
| PLEASE READ ALL INSTRUCTI application variety. Data for quar titative data. Comparative data sh- e.g., The Munsell Plant Tissue Colo | ntitative plant charac ould be determined (| ters should be based o | on a minimum of I | 00 plants. Include le | ading zeros when nec | cssary (c.g., 0 8 | 9) for quan- |
| 1, WINTERHARDINESS: | | | | | | | |
| - [] a | - Mari Mari Miland | d. (0)(5 104) | | | | | |
| 3 6 7 | 7] CLASS: 1 = Very Non-Winterhardy (CUF 101) 3 = Intermediately Non-Winterhardy (Mesilla) 5 = (Du Puits) 7 = (Ranger) 9 = Extremely Winterhardy (Norseman) | | | nardy (Moapa 69) hardy (Lahontan) Winterhardy (Saranac) (Vernal) | | | |
| T | EST LOCATION: | Johnston, I | A | | | | |
| 2. FALL DORMANCY: | FA | ALL DORMANCY (D | ETERMINED FRO | M SPACED PLANTI | INGS) | | |
| | | | | REGROWTH SCORE | OR AVERAGE HEIGH | <u> </u> | Τ |
| TESTING INSTITUTION | DATE OF | DATE REGROWTH | APPLICATION | | CHECK VARIETIES | | LSD .06 |
| AND LOCATION | LAST CUT | SCORED | VARIETY | Vernal | Ranger | Saranac | 1 |
| Pioneer Hi-Bred International, Inc. | 9/14/90 | 10/12/90 | 9.9 | 8.4 | 9.7 | 11.8 | 1.4 |
| Johnston, IA | | | | | | | |
| CUF 101, Moapa 69, Mesilia, Lahont | Du Buita Casasa I | <u> </u> | | | <u> </u> | 1 | <u> </u> |
| • | | | | | | | * |
| (, 2) | 4 | t height mea | asured in c | III s | | ······································ | - |
| Fall Growth Habit (Deter | mined from Fall Dorma | incy Trials) | | · | | • | |
| | = Erect (CUF 101) = Semidecumbent (Ver | | ierect (Mesilla) umbent (Norseman) | 5 = Intermediate | (Saranac) | | |
| | | | | | | | · · · · · · · · · · · · · · · · · · · |
| | st (CUF 101) | | : (Saranac) | 5 = Intermediate | s (Ranger) | 7 = Slow (Vernal) | |
| 9 = Very Sto | w (Norseman) | | | | | | |
| | | | | | | _, | |
| 4. AREAS OF ADAPTATION IN U.S. | (Where tested and prov | en adapted): | | | | | |
| Primary Area of Adaptation | on . | • | | 2 6 011 | ner Areas of Adaptation | | |
| | ** | | | | | | |
| 1 = North Co 5 = Moderate | intral ely Winterhardy (interm | 2 = East Central ountain | 3 = Soi 6 = Winterhardy Inte | utheast ermountain | 4 = Southwest 5 7 = Great Plains | 和国外 | 7 |
| 8 = Other (S | pecify) | | | | - 7 | SHU KE | |
| | | 2 to 1 to 2 | | | | | 4, |
| | | | | | | N | V |
| | | • • | • | ÷ | • | | |
| | | | *. | | <u> </u> | · | |
| 5. FLOWERING DATE (When 10% of) | | vers at time of first sprin | og cut): | | | | |
| Days Earlier Than | | | | • | | | |
| Same At | | 1 = CUF | 101 | 2 = Mesilla | 3 = Serenec 4 = | Vernal 5 = No | remen. |
| Days Later Than | TEST LOCATION: | | | | | | 4 |
| | | | | | | | |

| S'PLANT COLOR (Determin | ed from twelthy regrowth 3 | wroks after first sp | ring out, controlling | g ballourpers if necess | ıry) | | |
|--------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1 - Very Date G | reen (524) | 2 > Dark Green | (Vernal) | 3 / Leiht Green (| [Hangert | | 9100154 |
| | VALUE (Specify charriesed | | | | | — | 7100104 |
| APPLICATION | VARIETY: | | | | · · · · · · · · · · · · · · · · · · · | | |
| VERNAL: | | <u> </u> | | | | · · · · · · · · · · · · · · · · · · · | |
| 7. CROWN TYPE (Determin | | | | | | | |
| | - | | | | | | |
| Noncreeping | | (Vernat) | 2 * Intermediate i | Saranac) | 3 - Narrow f | CUF 101) | |
| Creeting Typ | Оссери | ng Rooted (Rangela | | 5 • Abizomatous | | | |
| 8. FLOWER COLOR (Deterr | nine frequency of plants for | each color class as | defined by USDA | Agriculturei Handbook | No. 424 (Bar | nes 1972), allowing al | plants in plot to flower): |
| 0 / 2 % Purple and | Violet (Subclasses 1,1 to 1,4 |) | 0 2 | ini | lasses 2.3 and | | |
| 0 0 8 % Variegated | Other Than Blue (Subclasses | 2.1, 2.2, 2.5 to 2.9 | , [| t % Yellow (Su | | | |
| % Cream (Clas | s 31 | | <u> </u> | | | , | • |
| TEST LOCAT | ION:Johnsto | on, IA | | t White (Clas | \$ 51 | | |
| 9. POD SHAPE (Determine (r | | ollowice and -t- | | | | | |
| | ed (One or more coils, cente | | | | | | · · · · · · · · · · · · · · · · · · · |
| | | r more or less close | d) | 1 3 Loosely Coi | iled (One or m | nare coils, center cons | picuously open) |
| % Sickle (Less | | | | TEST LOCAT | | | |
| 10. PEST RESISTANCE: Provinde: | rde in the appropriate colum Kiscores (ASI), least significa | in, trial data for app | olication variety, an | d resistant (R) and sus | ceptible (S) c | heck varieties, synthet | ic generation tested, everage se |
| •valç | ration. Describe scoring syst | iom, and any test no | Acadissa subject diss | | f test, year, ar | d location of test, and | ic generation tested, everage se I whether test is a field or tabo at data from other test years o |
| locat | ions should be presented wh | Minever available on | | O'A NOM STANDARD MARK | mous proposed | DY EIGIN (1982), Tri | at data from other test years o |
| 2070 | 5. Although competitions w | jarmpiasm lines liste rith check variaties (| id below can be ob listed below are pre | tained from the USDA | Field Crops I | Laboratory, Bidg. 001 | , Rm. 335, BARC-West, Beltsvi commended by Elgin (1982) n |
| A. DISEASE RESISTANCE: | nted. | · | | ,, | | | commended by Elgin (1982) n |
| THE PROCESS AND CE | 4 | r i | | | | | |
| DISEASE | VARIETY | SYN. GEN. | PERCENT RESISTANT | NUMBER OF | | ASI | INSTITUTION YEAR LOC |
| DISEASE | VARIETY | SYN. GEN. TESTED | PERCENT RESISTANT PLANTS | NUMBER OF PLANTS TESTED | ASI | ASI LSD .05 | INSTITUTION, YEAR, LOC FIELD OR LABORATO |
| Anthracnose, Race 1 | VARIETY Application R | | RESISTANT | Approx | ASI | LSD .05 | FIELD OR LABORATO |
| Anthracnose, Race 1 | | TESTED | RESISTANT PLANTS 40.7 | PLANTS TESTED | ASI | Percent | Pioneer Hi-Bre |
| Anthracnose, Race 1 | Application R | TESTED 1 | 40.7 65.0 | Approx 300 | ASI | LSD .05 | Pioneer Hi-Bre |
| Anthracnose, Race 1 | Application R Arc (B) Saranac AR | TESTED | 40.7 65.0 43.1 | Approx 300 | ASI | Percent Resistant | Pioneer Hi-Bre International, 1990 Johnston, IA |
| Anthracnose, Race 1 | Application R Arc (A) Saranac AR Saranac (S) | 1 (R) | 40.7 65.0 43.1 0.0 | Approx 300 | | Percent Resistant Plants 18.4 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory |
| Anthracnose, Race 1 | Application R Arc (A) Saranac AR Saranac (S) SCORING SYSTEM: P | 1 (R) ercent su | 40.7 65.0 43.1 0.0 | Approx 300 "" "" eedlings. | Data a | Percent Resistant Plants 18.4 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory |
| Anthracnose, Race 1 (Colletotrichum trifolii) | Application R Arc (B) Saranac AR | 1 (R) ercent su | 40.7 65.0 43.1 0.0 | Approx 300 "" "" eedlings. | Data a | Percent Resistant Plants 18.4 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory |
| Anthracnose, Race 1 | Application R Arc (A) Saranac AR Saranac (S) SCORING SYSTEM: P | 1 (R) ercent su | 40.7 65.0 43.1 0.0 rviving s ioneer Hi | Approx 300 "" "" eedlingsBred Inter | Data a | Percent Resistant Plants 18.4 djusted to | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (A) Saranac AR Saranac (S) SCORING SYSTEM: P resistant p1 | (R) ercent su ants by P | 40.7 65.0 43.1 0.0 | Approx 300 "" "" eedlingsBred Inter | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (A) Saranac AR Saranac (S) SCORING SYSTEM: P resistant p1 | (R) ercent su ants by P | 40.7 65.0 43.1 0.0 rviving s ioneer Hi | Approx 300 "" "" eedlingsBred Inter | Data a | Percent Resistant Plants 18.4 djusted to | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (R) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) | (R) ercent su ants by P | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 | Approx 300 "" "" eedlings. Bred Inter Approx 160 "" | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (R) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) | (R) ercent su ants by P | 40.7 65.0 43.1 0.0 rviving s ioneer Hi 11.1 55.0 2.4 | Approx 300 "" "eedlingsBred Inter Approx 160 "" | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (R) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P SCORING SYSTEM: P | (R) ercent su ants by P I | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving s | Approx 300 "" "" eedlings. Bred Inter Approx 160 "" eedlings. | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (R) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) | (R) ercent su ants by P I | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving s | Approx 300 "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 | Application R Arc (R) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P SCORING SYSTEM: P | (R) ercent su ants by P I | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving sby Pionee | Approx 300 "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I | Data a cnation | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (H) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Serenac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: | (R) ercent su ants by P I ercent su t plants | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving s | Approx 300 "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I | Data a | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc. ASI 0.98 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (H) Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Serenac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: | (R) ercent su ants by P I ercent su t plants | A0.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving sby Pionee | Approx 300 "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I | Data a cnation | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Serenec AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) | (R) ercent su ants by P I ercent su t plants | RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving soby Pionee 55.4 42.0 | eedlingsBred Inter Approx 160 "" eedlings. Thi-Bred I Approx 175 | Data a nation Data a nterna 5.94 4.87 | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P CORING SYSTEM: P Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Narraganiett (S) | (R) ercent su ants by P I ercent su t plants | 85.51ANT PLANTS 40.7 65.0 43.1 0.0 rviving s ioneer Hi 11.1 55.0 2.4 rviving s by Pionee 55.4 42.0 3.0 | eedlingsBred Inter Approx 160 "" eedlings. Thi-Bred I Approx 175 "" | Data a nterna 5.94 4.87 2.22 | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Nerresponsett (S) SCORING SYSTEM: P1 | ercent su ants by P I ercent su ants by I lants score | 85.51ANT PLANTS 40.7 65.0 43.1 0.0 rviving s ioneer Hi 11.1 55.0 2.4 rviving s by Pionee 55.4 42.0 3.0 ced 7-9 (6 | eedlingsBred Inter Approx 160 "" eedlings. Thi-Bred I Approx 175 "" on a 1-9 sc | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) Bacterial Wilt (Corynebacterium insidiosum) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan Application HR Vernal (R) Narraganizati (S) SCORING SYSTEM: P p1ant) consider | ercent su ants by P I ercent su t plants lants scondered residered | ## RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving siby Pionee 55.4 42.0 3.0 ced 7-9 (distant.) | Approx 300 "" "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I Approx 175 "" on a 1-9 sc Data adjust | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, F Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Nerresponsett (S) SCORING SYSTEM: P1 | ercent su ants by P I ercent su t plants lants scondered residered | ## RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving siby Pionee 55.4 42.0 3.0 ced 7-9 (distant.) | Approx 300 "" "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I Approx 175 "" on a 1-9 sc Data adjust | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) Bacterial Wilt (Corynebacterium insidiosum) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Narraganisati (S) SCORING SYSTEM: P1 plant) consider by Pioneer III | ercent su ants by P I ercent su t plants lants scondered residered | ## RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving siby Pionee 55.4 42.0 3.0 ced 7-9 (distant.) | Approx 300 "" "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I Approx 175 "" on a 1-9 sc Data adjust | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) Bacterial Wilt (Corynebacterium insidiosum) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Narraganisati (S) SCORING SYSTEM: P1 plant) consider by Pioneer III | ercent su ants by P I ercent su t plants lants scondered residered | ## RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving siby Pionee 55.4 42.0 3.0 ced 7-9 (distant.) | Approx 300 "" "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I Approx 175 "" on a 1-9 sc Data adjust | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI Field |
| Anthracnose, Race 1 (Colletotrichum trifolii) Anthracnose, Race 2 (Collectotrichum trifolii) Bacterial Wilt (Corynebacterium insidiosum) | Application R Arc (R) Saranac AR Saranac AR Saranac (S) SCORING SYSTEM: P Tesistant p1 Application LR Saranac AR (R) Arc (S) SCORING SYSTEM: P 55% resistan: Application HR Vernal (R) Necrepansett (S) SCORING SYSTEM: P plant) consider positions of the position of the positions of the position of the positions of the positions of the position of the position of the positions of the position | ercent su ants by P I ercent su t plants lants scondered residered | ## RESISTANT PLANTS 40.7 65.0 43.1 0.0 rviving sioneer Hi 11.1 55.0 2.4 rviving siby Pionee 55.4 42.0 3.0 ced 7-9 (distant.) | Approx 300 "" "" eedlingsBred Inter Approx 160 "" eedlings. r Hi-Bred I Approx 175 "" on a 1-9 sc Data adjust | Data a nterna 5.94 4.87 2.22 ale wh | Percent Resistant Plants 18.4 djusted to al, Inc. Percent Resistant Plants 14.5 djusted to tional, Inc ASI 0.98 Percent Resistant Plants 14.1 ere 9=no di | Pioneer Hi-Bre International, 1990 Johnston, IA Laboratory Arc at 65% Pioneer Hi-Bre International, 1990 Quarryville, P Laboratory Saranac AR at Pioneer Hi-Bre International, 1990 Arlington, WI |

| DISEASE | VARIETY | SYN. GEN. TESTED | PERCENT RESISTANT PLANTS | NUMBER OF PLANTS TESTED | ASI | ASI LSD .05 | INSTITUTION, YEAR, LOCATION, FIELD OR LABORATORY |
|----------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|--------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| Downy Mildew (Peronospora trifoliorum) | Application | | | | | | 9100154 |
| Isolete, if known: | Saranac (R) | | | | | 1 | |
| <u></u> | Kanza (S) | | | | | | |
| · · · · · · · · · · · · · · · · · · · | SCORING SYSTEM: | | .1 | | | | L |
| Fuserium Wilt Fuserium oxysporum . mediceginis) | Application HR | 1 | 61.0 | Approx . 225 | 1.89 | ASI 0.60 | University of |
| | Agate (R) | | 54.0 76.7 | 11 | 2.40 1.56 | Percent Resistant | Minnesota 1990 |
| • | NarragansettX(X) .(M | R) | 28.9 | 11 | 3.48 | Plants 14.08 | Rosemount, MN Field |
| | MNCN-1 (S) scoring system: 5=dead plan | Plants s | cored 0 ar | d 1 (on a | 1 4 58 0-5 sca | le, where | 0=no disease and |
| Phytophthora Root Rot | iplants by t | he Unive | rsity of M | innesota | ta adju T | ASI 0.59 | ate at 54% resista University of |
| (Phytophthora megasperma 1, medicaginis) | Application HR | 1 | 52.6 | Approx 225 | 3.30 | Percent | Minnesota |
| | Agate (R) | | 43.0 | 11 | 3.29 | Resistant Plants | 1990 St. Paul, MN |
| | Saranac (S) | | 1.7 | 11 | 5.28 | 16.39 | Field |
| | SCORING SYSTEM: P | lants sco | ored 1 and | 2 (on a 1- | -6 scal | e, where 1 | =no disease and 6= |
| Verticillium Wilt | plants by t | conside | red resist rsity of M | ant. Data | adjust | | e at 43% resistant |
| verticillium villt [Verticillium elboetrum] | Application HR | ie unive | 67.0 | Approx | 2.77 | ASI 0.69 Percent | University of Wisconsin |
| | Vertus (A) | | 40.0 | 150 " | 3.13 | Resistant Plants | 1990 Madison, WI |
| | t | | | | | | |
| | Saranac (S) | | 11.0 | . 11 | 4.51 | 23.0 | Laboratory |
| | SCORING SYSTEM: | Plants so | cored 1 (o | n a 1-5 sca | l ale, who | ere 1=no d | Laboratory isease and 5=dead |
| Other (Specify) | scoring system: plant) cons | idered re | cored 1 (oesistant. | n a 1-5 sca Data adjus | l ale, who | ere l=no d Vertus at | Laboratory isease and 5=dead 40% resistant pla |
| • • • | SCORING SYSTEM: | idered re | cored 1 (oesistant. | n a 1-5 sca Data adjus | l ale, who | ere 1=no d Vertus at ASI 0.70 Percent | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In |
| | scoring system: plant) cons | idered re | cored l (o esistant. Wisconsi | n a 1-5 sca Data adjus n. Approx | ale, who | ere 1=no d Vertus at ASI 0.70 | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 |
| | scoring system: plant) cons: by the University HR (R) WAPH-1 | idered re | cored 1 (oesistant. Wisconsi 65.0 50.0 | n a 1-5 sca Data adjus h. Approx 185 | 6.17 5.56 | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory |
| | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: | rsity of | cored 1 (oesistant. Wisconsi 65.0 50.0 2.0 cored /-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 s | ale, who sted to 6.17 5.56 1.68 ccale, v | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= |
| Aphanomyces | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: | rsity of | cored 1 (o esistant. Wisconsi 65.0 50.0 2.0 cored 7-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 sant. Data | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory |
| Aphanomyces | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: dead plant) plants by Pi | rsity of | cored 1 (o esistant. Wisconsi 65.0 50.0 2.0 cored 7-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 sant. Data | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= |
| Aphanomyces | scoring system: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: 1 dead plant) plants by Paricestion | rsity of | cored 1 (o esistant. Wisconsi 65.0 50.0 2.0 cored 7-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 sant. Data | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= |
| Aphanomyces | SCORING SYSTEM: plant) cons: by the University Application HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: dead plant) plants by Prapplication (R) | rsity of | cored 1 (o esistant. Wisconsi 65.0 50.0 2.0 cored 7-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 sant. Data | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= |
| Other (Specify) | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: 1 dead plant) plants by Prapplication (R) (S) | l ered resity of | cored 1 (oresistant. Wisconsi 65.0 50.0 2.0 cored /-9 red resistBred Int | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 s ant. Data ernational, | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no ed to WAPH- | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= -1 at 50% resistan |
| Other (Specify) | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: 1 dead plant) plants by Prapplication (R) (S) | rsity of | cored 1 (o esistant. Wisconsi 65.0 50.0 2.0 cored 7-9 | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 sant. Data | 6.17 5.56 1.68 scale, valuete | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= |
| Other (Specify) NSECT RESISTANCE: INSECT | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: dead plant) plants by Praphication (R) (S) SCORING SYSTEM: | rsity of lants so consider oneer Hi | cored 1 (oesistant. Wisconsi 65.0 50.0 2.0 cored /-9 ced resistBred Int | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 s ant. Data ernational, | 6.17 5.56 1.68 scale, valuate | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no ed to WAPH- | isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and I= -1 at 50% resistan |
| Other (Specify) Aphanomyces Other (Specify) INSECT RESISTANCE: INSECT Alfalia Weevil (Hypera postica) | SCORING SYSTEM: plant) cons: by the University Application HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: dead plant) plants by Practication (R) (S) SCORING SYSTEM: VARIETY | rsity of lants so consider oneer Hi | cored 1 (oesistant. Wisconsi 65.0 50.0 2.0 cored /-9 ced resistBred Int | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 s ant. Data ernational, | ale, who sted to 6.17 5.56 1.68 scale, wadjuste Inc. | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no ed to WAPH- | Laboratory isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and 1= -1 at 50% resistan |
| Other (Specify) INSECT RESISTANCE: INSECT | SCORING SYSTEM: plant) cons: by the University HR (R) WAPH-1 (S) Saranac SCORING SYSTEM: dead plant) plants by Prapplication (R) (S) SCORING SYSTEM: VARIETY | rsity of lants so consider oneer Hi | cored 1 (oesistant. Wisconsi 65.0 50.0 2.0 cored /-9 ced resistBred Int | n a 1-5 sca Data adjus n. Approx 185 " (on a 1-9 s ant. Data ernational, Defoliation in PERCENT OF RESISTANT CHECK | ale, who sted to 6.17 5.56 1.68 scale, wadjuste Inc. | ere 1=no d Vertus at ASI 0.70 Percent Resistant Plants 6.0 where 9=no ed to WAPH- | isease and 5=dead 40% resistant pla Pioneer Hi-Bred International, In 1990 Arlington, WI Laboratory symptoms and I= -1 at 50% resistan |

| | · 1 | l l | | | | - | |
|----------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------|--------------------------------------------------|--------------------------------------------|--------|-------------------------|--------------------------------------------------------------------------------------------------|
| INSECT | VARIETY | SYN, GEN. TESTED | PERCENT SEEDLING SURVIVAL | NUMBER OF SEEDLINGS TESTED | ASI | ASI LSD .05 | INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY |
| Blue Alfalfa Aphid (Acyrthosiphon kandai) | Application | | · | | | | |
| e e | CUF 101 (R) | | | | | - | |
| | PA-1 (S) | | | | | | |
| | SCORING SYSTEM: | | | | - | | |
| Pea Aphid (Acyrthosiphan pisum) | Application | | | | | | |
| | Konza (R) | | | | | | |
| | Ranger (S) | | | | | | |
| | SCORING SYSTEM: | | | | | | |
| Spotted Alfalfa Aphid (Therioaphis maculata) | Application R | 1 | 39.1 | Approx 200 | 2.94 | ASI 1.0 Percent | Pioneer Hi-Bred International, In |
| Biotype, if known: | Kanza (R) | | 70.0 | 11 | 4.01 | Resistant Plants | 1990 Kerman, CA |
| | Ranger (S) | · - | 3.5 | 11 | 1.61 | 27.7 | Laboratory |
| | | | | / | | 1 0 | |
| | scoring system: plant) cons | | | | | | |
| INSECT | plant) cons | idered re | | Data adjus | | | 0% resistant plant |
| INSECT Potato Leafhopper Yellowing (Empossca fabse) | plant) cons | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing | plant) cons | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing | plant) cons by Pioneer VARIETY Application | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | symptoms and l=dea. % resistant plant INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY |
| Potato Leafhopper Yellowing | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing (Empossca fabse) | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing (Empossca fabse) | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: Application | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing (Empossca fabse) | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: Application (R) | idered re | ntermatio | Data adjus | ted to | Kanza at 7 | 0% resistant plant |
| Potato Leafhopper Yellowing (Empossca fabse) | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: Application (R) (S) | idered re | PERCENT RESISTANT | Data adjus | ted to | Kanza at 7 | 0% resistant plant INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY |
| Potato Leafhopper Yellowing (Emposca fabse) Other (Specify) NEMATODE RESISTANCE: | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: (R) (S) SCORING SYSTEM: | Hi-Bred restant syn. GEN. TESTED | esistant. ntermetio RESISTANT PLANTS PERCENT | Data adjus nal Inc number of plants tested | ted to | Kanza at 7 ASI LSD .05 | 0% resistant plant INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY INSTITUTION, YEAR, LOCATION, |
| Potato Leafhopper Yellowing (Emposca fabse) Other (Specify) NEMATODE RESISTANCE: NEMATODE Northern Root Knot | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: Application (R) (S) SCORING SYSTEM: | Hi-Bred restant syn. GEN. TESTED | PERCENT RESISTANT | Data adjus nal Inc number of plants tested | ted to | Kanza at 7 ASI LSD .05 | 0% resistant plant INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY INSTITUTION, YEAR, LOCATION, |
| Potato Leafhopper Yellowing (Emposca fabse) Other (Specify) NEMATODE RESISTANCE: NEMATODE Northern Root Knot | plant) cons by Pioneer VARIETY Application MSA-CW3An3 (R) Ranger (S) SCORING SYSTEM: Application (R) (S) SCORING SYSTEM: VARIETY Application | Hi-Bred restant syn. GEN. TESTED | PERCENT RESISTANT | Data adjus nal Inc number of plants tested | ted to | Kanza at 7 ASI LSD .05 | 0% resistant plant INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY INSTITUTION, YEAR, LOCATION, |

| NEMATODE | VARIETY | SYN. GEN. TESTED | PERCENT RESISTANT PLANTS | NUMBER OF PLANTS TESTED | ASI | AŞI LSD .05 | INSTITUTION, YEAR, LOCATION FIELD OR LABORATORY |
|-----------------------------------------------|-----------------|---------------------|--------------------------------|----------------------------|------|---------------------|----------------------------------------------------|
| Southern Root Knot (Melaidagyne incognite) | Application | | | | | | 910015 |
| | Mosps 69 (R) | | | | | | |
| | Lahonten (S) | | | | | | |
| | SCORING SYSTEM: | | | | | | |
| Stem Nematode (Ditylenchus dipseci) | Application LR | 1 | 8.5 | Approx 200 | 1.77 | ASI 0.45 Percent | Pioneer Hi-Bred International, Ir |
| | Lahontan (R) | | 50.0 | 11 | 3.26 | Resistant | · |
| ÷ | Renger (S) | | 4.3 | ft | 1.50 | 11.4 | Laboratory |
| | • | | | | - | | symptoms and l=dea t 50% resistant |
| Other (Specify) | plants by Pi | | | | | | |
| | (R) | | | | | | |
| <u> </u> | (5) | | | | | 1 | · |

11. INDICATE THE VARIETY THAT MOST CLOSELY RESEMBLES THE APPLICATION VARIETY FOR EACH OF THE FOLLOWING CHARACTERS:

| CHARACTER | VARIETY | CHARACTER | VARIETY |
|------------------------|---------|-----------------------------|---------|
| Winterhardiness | Ranger | Plant Color | - |
| Recovery After 1st Cut | Ranger | Crown Type | - |
| Area of Adaptation | 532 | Combined Disease Resistance | Dart |
| Flowering Date | _ | Combined Insect Resistance | 5432 |

REFERENCES

Barnes, D.K. 1972. A System for Visually Classifying Alfalfa Flower Color. U.S. Dep. Agric. Handb. 424. 18 pp. (Note: Greenish cast of plate 6, A and B is an artifact of printing, actual colors a blend of yellow and white.)

Elgin, J.H., Jr., (ed.). 1982. Standard Tests to Characterize Pest Resistance in Alfalfa Cultivars. U.S. Dep. Agric, Tech. Bull. (In Press).

Gunn, C.R., W.H. Skrdla, and H.C. Spencer. 1978. Classification of Medicago sativa L. using legume characters and flower colors. U.S. Dep. Agric. Tech. Bull. 1574. 84 pp.

Munsell Color Co. 1977. Munsell Plant Tissue Color Charts. Munsell Color Co., Inc. Baltimore.

NOTE: Any additional descriptive information and supporting documentation may be provided as Exhibit D.

EXHIBIT D

152521

- 1. 5252 is a synthetic variety with 197 parent plants originating from an experimental line tracing to the varieties 5373 and Surpass. Parent plants were selected through phenotypic recurrent selection for resistance to Phytophthora root rot and/or Aphanomyces root rot. Germplasm sources are M. falcata (8%), Ladak (8%), M. varia (23%), Turkistan (10%), Flemish (42%), Chilean (7%), and Peruvian (2%).
- 2. 5252 is adapted to and intended for use in the central and northern region of the United States for hay, haylage, greenchop, and dehydration. The states in which 5252 is being tested are Iowa, Illinois, Minnesota, New York, Pennsylvania, Wisconsin, Oregon, and Washington. The variety has also been tested using experimental designation YAM93.
- 3. 5252 is a dormant cultivar with fall dormancy similar to Ranger. Flower color in the Syn 1 generation is approximately 72% purple and 28% variegated with traces of yellow, white, and cream. Growth habit is erect in midsummer and semi-erect in the fall.
- 4. 5252 has high resistance to Aphanomyces (Race 1), bacterial wilt, Fusarium wilt, Phytophthora root rot, and Verticillium wilt; resistance to anthracnose (race 1) and spotted alfalfa aphid; low resistance to anthracnose (race 2). YAM93 has not been adequately tested for pea aphid, blue alfalfa aphid or stem nematode.
- 5. Breeder seed (Syn 1) was produced in one year on parent plants in "cage isolation". Seed classes will be breeder, foundation (Syn 2 or Syn 3) and certified (Syn 2, Syn 3, or Syn 4). Foundation seed may be produced from breeder or foundation. The second generation foundation may be produced at the discretion of Pioneer Hi-Bred International Inc. Limitations on ages of stand will be three years and five years, respectively, for foundation seed and certified seed. Sufficient breeder and foundation seed for the projected life of the variety will be maintained by Pioneer Hi-Bred International, Inc.
- Seed will be marketed in the spring of 1992.
- Application for Plant Variety Protection will be made, and the certification option not will be requested.

EXHIBIT E

STATEMENT OF THE BASIS OF APPLICANT'S OWNERSHIP

Pioneer Hi-Bred International, Inc., Des Moines, Iowa, is the employer of the plant breeders involved in the development and evaluation of 5252. Pioneer Hi-Bred International, Inc. has the sole rights and ownership of 5252.